

THE DIFFERENTIATED WEED CONTROL METHODS IN ONION CROPS ON THE CAMBIC CHERNOZEM IN SOUTH ROMANIA

Mihai GÎDEA¹

¹University of Agronomic Sciences and Veterinary Medicine, Romania
Corresponding author: Mihai GÎDEA, E-mail: gideam@yahoo.com

Received November 26, 2012

By limiting the access to vegetation factors, weeds determine significant losses in all agricultural crops, and, depending on the weed coverage they can result in crop failure. That's why there can be no crop technologies that don't take into account weed control methods. This research aimed to compare the large-scale weed control methods used on the entire surface with the precise application of herbicides according to the local weed species. The findings were that the precise application of weed control methods determined a significant reduction in weed coverage as well as a reduction of production costs in comparison with the uniform application of herbicides on the entire surface.

Key words: onion, weed management, weed mapping, weed control.

INTRODUCTION

The precise agricultural system, as the most efficient system of sustainable agriculture, equals the application of technological measures according to the specific features of each lot.

The biology of onion plants and their slow growth in the first 6 to 8 weeks after emergence results in a low capacity of competing with weeds, which offers weeds an edge when using vegetation factors.

This research aimed to assess the ability to decrease herbicide use in onion crops by applying precise herbicides instead of the classical uniform application on the entire surface.

The starting assumption of the research was that weeds aren't uniformly spread in a crop and anyway weed coverage is under 100%.

MATERIALS AND METHODS

In order to properly assess the effectiveness of a weed control method, it's necessary to determine weed coverage before and after applying the respective method.

Weed control in the precise agricultural system is based on the differentiated application of weed control methods according to weed mapping and not on measures applied uniformly on the entire surface.

The research was carried out in a monofactorial experiment, on the cambic chernozem in south Romania. (Coordinates Stereo 70).

The experimental variants were V1 uniformly applied with herbicides for controlling dicotyledonous and monocotyledonous weeds; V2 applied with herbicides according to the weed species locally identified.

In order to control the dicotyledonous species Lontrel 300 0.3/ha (clopyralid 300 g/l) was used, and Fusilade Super 1.5l/ha (fluazifop-P-butyl 150 g/l) was used to control monocotyledonous species.

The surface of the experimental lots was of approximately 1000 sqm, in 3 repetitions.

The coordinates of the experimental lots were determined with a GPS RTK Trimble Ag 262 in a WGS 84 system. The field data were processed in Autocad and correlated with the zone map. TopoLT was used to generate a point network of 2 m × 2 m. Then the coordinates of the network points were loaded in a GPS. The measurements for the weed mapping were performed only in the network knots by using the 0,25 sqm (50 × 50 cm) frame.

The average weed occurrence, prevalence and persistence were calculated for each of the two experimental variants, before and after herbicide application.

The average $a = S/N$, where S is the total number of plant individuals belonging to a certain species found in all measurement points, and N is the number of measurement points and “ a ” is the mean number of weed plants belonging to a certain species per sqm.

The participation (prevalence) $P\% = m*100/M$, where $M = \sum m$ is the mean number of weed plants/sqm, the sum of all averages of weed species identified.

The persistence, $K\% = n*100/N$, where n is the number of points where a certain species occurred.

By using the gathered data the distribution maps for the monocotyledonous and dicotyledonous species were drawn for all 6 lots (2 variants \times 3 repetitions). By using these maps the herbicides from the second variant were applied differently according to the distribution zones of the identified weed species.

RESULTS

The weed mapping before applying herbicides revealed the occurrence of 8 annual dicotyledonous species: *Amaranthus retroflexus*, *Xanthium strumarium*, *Sinapis arvensis*, *Poligonum convolvulus*, *Chenopodium album*, *Galinsoga parviflora*, *Vicia angustifolia* and 2 perennial dicotyledonous species *Convolvulus arvensis*, *Cirsium arvensis*; 2 annual monocotyledonous species *Setaria sp.*, *Echinochloa cruss galli* and a perennial monocotyledonous species *Sorgum halepense*.

The data regarding the average weed occurrence, prevalence and persistence before herbicide application are gathered in Table 1.

The analysis of the data regarding the mean number of weed plants reveals that: the mean number of weed plants per sqm varied from 58.5 and 57.6; the number of annual dicotyledonous weed plants was between 26 and 26.2 plants/sqm; and the number of annual monocotyledonous weed plants was between 29.5 and 28.4 plants/sqm.

As far as the weed species prevalence is concerned, the highest percentage was of annual monocotyledonous species, with over 50.4%, and lowest percentage was of perennial monocotyledonous species, with less than 1.3%.

As far as the weed species persistence is concerned, the highest percentage was of annual monocotyledonous species, with over 50.4%, and lowest percentage was of perennial monocotyledonous species, with less than 1.3%.

The persistence of weed species in the measurement points was lower than 61.2%.

The comparative analysis between the data recorded in the two experimental variants, before applying the postemergent treatments, showed there were no significant differences regarding the number of weed species, the mean number of weed plants per sqm, the prevalence and the persistence.

Table 1

Weed mapping of experimental lots before herbicide application

Species	Variant 1			Variant 2		
	M	P	C	M	P	C
<i>Amaranthus retroflexus</i>	5.5	9.4	48.3	5.8	10.1	54
<i>Xanthium strumarium</i>	2.6	4.4	36,2	2.8	4.9	50.4
<i>Vicia angustifolia</i>	3.8	6.5	18,7	3.3	5.7	12,8
<i>Solanum nigrum</i>	2.8	4.8	43,1	2.7	4.7	48
<i>Poligonum convolvulus</i>	3.8	6.5	36.7	4.1	7.1	49.7
<i>Chenopodium album</i>	1.1	1.9	34.9	1.4	2.4	57.9
<i>Sinapis arvensis</i>	2.8	4.8	22,9	2.3	4.0	19.7
<i>Galinsoga parviflora</i>	3.6	6.2	61.2	3.8	6.6	60
Annual dicotyledonous	26	44.4		26.2	45.5	
<i>Convolvulus arvensis</i>	0.5	0.9	20.6	0.3	0.5	37.7
<i>Cirsium arvensis</i>	1.2	2.1	26.7	1.3	2.3	32
Perennial dicotyledonous	1.7	2.9		1.6	2.8	
<i>Setaria sp.</i>	16.9	28.9	41.1	17.2	29.9	42.3
<i>Echinochloa cruss galli</i>	12.6	21.5	37.4	11.2	19.4	60.2
Annual monocotylenous	29.5	50.4		28.4	49.3	
<i>Sorgum halepense</i>	1.3	2.2	11.3	1.4	2.4	20.2
Perennial monocotylenous	1.3	2.2		1.4	2.4	
Total	58.5	100		57.6	100	

According to these data the distribution maps with the monocotyledonous and dicotyledonous weed populations were drawn. In variant no. 1 the lots were uniformly applied with the two herbicides Lontrel 300 and Fusilade Super, and in variant no. 2 the treatments were applied differently, according to the distribution zones of the monocotyledonous and dicotyledonous populations.

The analysis of the mean data regarding weed coverage shows that, after applying herbicides in both studied variants, the average number of weed plants in each biological group was lower. The weed control extent was 89.9% when applying the two herbicides uniformly on the entire surface and 86.8% when applying the two herbicides locally, only where weed plants occurred.

When calculating the surfaces actually treated with herbicides, the results showed that the surface actually treated with herbicides was of 68.3%, thus the recorded reduction of herbicide consumption was of 31.7%.

CONCLUSIONS


Before applying the herbicide treatments, the weed occurrence, expressed through the mean number number of weed plants, the prevalence of weed species and their persistence weren't significantly different for the two experimental variants.

Table 2

Comparative evaluation of weednes level before and after treatment application

Biological group	Variant 1				Variant 2			
	Average		Prevalence	Control extent	average		Prevalence	Control extent
	Before	After			Before	After		
Annual dicotyledonous	26	2.9	49.2	88.8	26.2	3.1	52.9	88.2
Perennial dicotyledonous	1.7	1.4	23.7	17.6	1.6	1.4	9.8	12.5
Annual monocotyledonous	29.5	1.0	16.9	96.6	28.4	2.2	19.6	92.3
Perennial monocotyledonous	1.3	0.6	10.2	53.8	1.4	0.9	17.6	35.7
Total	58.5	5.9	100.0	89.9	57.6	7.6	100.0	86.8

Figure 1. Biological group distribution before and after treatment





The application of herbicides resulted in a reduction of the total number of weed plants in both experimental variants.

The weed control extent were lower in the precise local application of herbicides than in the uniform application of the two herbicides on the entire surface.

The surface treated with herbicides reduced with 31.7% in the case of the precise local application, resulting in a corresponding reduction, of 31.7%, of the amount of herbicides used.

REFERENCES

1. Baghestani MA, Zand E, Soufizadeh S, Beygi MA. ,Evaluation of different empirical models of crop/weed competition to estimate yield and LAI losses from common lambsquarters (*Chenopodium album* L.) in maize (*Zea mays* L.). Pak J Biol Sci. 2007; 1,10(21): 3752-61.
2. Gail G. and co, Evaluating the potential for Site –specific Herbicide Application in soybean, weed technology, 2004, 18, 1101–1110.